



UNIVERSITI PUTRA MALAYSIA

**BASAL STEM ROT OF OIL PALM :
SCREENING SYSTEM AND BIOEFFICACY OF STEROL
BIOSYNTHESIS INHIBITORS AGAINST GANODERMA BONINENSE**

TEH KIM SING

FP 1996 14

**BASAL STEM ROT OF OIL PALM:
SCREENING SYSTEM AND BIOEFFICACY OF STEROL
BIOSYNTHESIS INHIBITORS AGAINST *Ganoderma boninense***

**By
TEH KIM SING**

**Thesis Submitted in Fulfilment of the Requirements
for the Degree of Master of Agricultural
Science in the Faculty of Agriculture
Unviversiti Pertanian Malaysia**

May 1996



ACKNOWLEDGEMENTS

The author expresses his sincere and profound gratitude to Associate Professor Dr. Sariah Meon, chairperson of the supervisory committee, members of the supervisory committee, Professor Dr. Abdul Rahman Abdul Razak, for their guidance, invaluable advice and patience throughout the duration of this study and in the preparation of this thesis; and Dr. Mohd. Zakaria Hussin for his useful comments. To Dr. Dieter Nordmeyer from Ciba the author is greatly indebted for his guidance, suggestions, constructive comments, constant encouragement and support, especially during the early part of the study.

Appreciation is also extended to the Ciba Research Station for providing facilities for field trials and for supplying the test fungicides, and to the Highland Reseach Unit for supplying the DxP oil palm seedlings for the trials. Special thanks to all the staff of Ciba, for their support and co-operation in the field works, in particularly the Disease Control Unit, Mr. Halim Baharol and Zulkifli for their excellent assistance and support.

The author also wishes to thank Miss Rose Lee for proof- read the final draft of the thesis; all laboratory staff of the Plant Pathology Department and staff of the Graduate School for their kind assistance.

The author owes so much to his wife Gait Sem, daughter Yuan-ning and son Qingyin for their love, understanding and patience. Last but not least, the author wishes to thank all those who in one way or another helped in these studies.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	ii
LIST OF TABLES	vi
LIST OF FIGURES	viii
LIST OF PLATES	ix
ABSTRACT	x
ABSTRAK	xii
CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	4
Status of <i>Ganoderma</i> spp. as a Pathogen of Basal Stem Rot (BSR) of Oil Palm	4
Symptoms of Basal Stem Rot on Oil Palm	6
Foliar Symptoms	6
Basidioma	7
Symptoms in the Roots and Stem Region (Bole) ...	8
Establishment of <i>Ganoderma</i> Infection by Inoculation	9
Management of Basal Stem Rot	11
Current Practises in Plantations	11
Potential Control Methods	13



	Page
Sterol Biosynthesis Inhibitor fungicides	15
Mode of Action	16
Chemical Grouping	17
Systemic Properties	17
Vapour Phase Activity	18
Fungicidal Activity	18
Non-target Effects	19
III MATERIALS AND METHODS	20
Screening System for Basal Stem Rot of Oil Palm	20
Source of Inoculum	20
Inoculum Preparation	20
Development of Screening Technique on Oil Palm Seedlings	21
Development of Screening Technique on Immature Oil Palm in the Field	24
Bioefficacy of Sterol Biosynthesis Inhibitors on <i>G. boninense</i>	24
<i>In vitro</i> Fungicide Tests	25
Bioassay of Fungicides	26
Curative Activity of the Triazole Fungicides	27

	Page
IV RESULTS	30
Screening System for Basal Stem Rot of Oil Palm	30
Development of Screening Technique on Oil Palm Seedlings	30
Development of Screening Technique on Immature Oil Palm in the Field	41
Bioefficacy of Sterol Biosynthesis Inhibitors on <i>G. boninense</i>	47
<i>In vitro</i> Fungicides Test	47
Bioassay of Fungicides	52
Curative Activity of the Triazole Fungicides	54
V DISCUSSION	58
Screening System for Basal Stem Rot of Oil Palm	58
Bioefficacy of Sterol Biosynthesis Inhibitors on <i>G. boninense</i>	61
VI SUMMARY AND CONCLUSION	65
LITERATURE CITED	69
APPENDICES	76
A Additional Tables	77
B Additional Figure	80
VITA	81

LIST OF TABLES

Table	Page
1 List of Triazoles and Its Dosage Used in the Curative Activity Experiments Against BSR of Inoculated Oil Palm Seedlings	28
2 Development of Basidioma of <i>Ganoderma</i> on Inoculated Oil Palm Seedlings from 30-162 Days After Inoculation (DAI)	37
3 Intensity and Incidence of <i>Ganoderma</i> Infection on Various Parts of the Inoculated Oil Palm Seedlings	41
4 Relationship between Various Infected Parts of Oil Palm Seedlings	41
5 Development of Basal Stem Rot-associated Symptoms on Inoculated Palms in the Field	45
6 Effect of Fungicides on Mycelial Growth of <i>G. boninense</i> on MEA	47
7 Effect of the Vapour Phase Activity of Morpholines and Triazoles on Mycelial Growth of <i>G. boninense</i>	48
8 Activity of Penconazole, Propiconazole and Difenoconazole on the Mycelial Growth of <i>G. boninense</i> with Time	52
9 Distributional Activity of Penconazole, Propiconazole and Difenoconazole in the Bole of Oil Palm Seedlings on the Mycelial Growth of <i>G. boninense</i> with Time.....	54
10 Effect of Triazoles on the Development of BSR by <i>G. boninense</i> on Inoculated Oil Palm Seedlings	56
11 Some Important Physical Parameters of the SBI fungicides Used in the Study	77
12 Analysis of Variance for Severity of Foliar Symptoms between Seedlings With and Without Friuting-bodies	78
13 Analysis of Variance for Relationship between Foliar Symptoms Against Bole and Root Infection of Inoculated Seedlings	78



Table		Page
14	Analysis of Variance for Relationship between Disease Penetration Against Bole and Root Infection of Inoculated Seedlings	78
15	Analysis of Variance for Relationship between Disease of Non-inoculated Root Against Bole and Inoculated Root Infection of Inoculated Seedlings	79

LIST OF FIGURES

Figure	Page
1 Diagrammatic Representations of the Bioassay Test on the Bole Tissues	27
2 Distribution of <i>Ganoderma</i> -associated Foliar Symptoms on 3 MAI Oil Palm Seedlings	31
3 Progressive Development of <i>Ganoderma</i> -associated Foliar Symptoms on Inoculated Oil Palm Seedlings	38
4 Distribution of <i>Ganoderma</i> -associated Symptoms on Various Parts of Inoculated Oil Palm Seedlings	40
5 Relationship between Severity of Foliar Symptoms and Growth of Inoculated Oil Palm Seedlings	42
6 Relationship between Root Infection of Inoculated Oil Palm Seedlings and Percent of Bole Infection	42
7 <i>In vitro</i> Bioefficacy of Morpholine Fungicides on Mycelial Growth of <i>G. boninense</i>	49
8 <i>In vitro</i> Bioefficacy of Triazole Fungicides on Mycelial Growth of <i>G. boninense</i>	50
9 <i>In vitro</i> Vapour Phase Activity of Fungicides on Mycelial Growth of <i>G. boninense</i>	53
10 Effect of Triazole Fungicides Against Basal Stem Rot on Inoculated Oil Palm Seedlings (Soil Drenched, 50 ml Fungicide Solution per seedling)	56
11 Diagrammatic Representations of the Cross Sections of Oil Palm Seedling	80

LIST OF PLATES

Plate	Page
1 Plastic Container Used in the Screening Technique	22
2 Inoculation Procedure Using the Split-Root Technique	22
3 Application of Fungicide Solution on the Non-inoculated Site	29
4 Comparative Response between Infected and Non-infected Roots of Oil Palm Seedlings	32
5 Formation of Thick Masses of Brownish White Mycelium Mat on the Oil Palm Seedlings at the Early and Late Stage of Colonisation	34
6 Fomation of White Button-like Basidioma Primodia at the Bole-root Junction of the Inoculated Seedling	35
7 A Typical Symptom Manifestation of the Bole Region of an Infected Seedling	36
8 Sequential Development of Foliar Symptoms on Infected Seedlings	39
9 Symptomatology of Infected Bole of the Inoculated Immature Oil Palm in the Field	44
10 Associated BSR Symptoms on Inoculated Palms	46
11 Vapour Phase Activity of Morpholines and Triazoles at 2000 ppm a.i.	51

Abstract of thesis submitted to the Senate of Universiti Pertanian Malaysia in fulfilment of the requirements for the degree of Master of Agricultural Science.

**BASAL STEM ROT OF OIL PALM: SCREENING SYSTEM AND
BIOEFFICACY OF STEROL BIOSYNTHESIS INHIBITORS
AGAINST *Ganoderma boninense***

By

TEH KIM SING

May 1996

Chairperson : Associate Professor Dr. Sariah Meon

Faculty : Agriculture

Studies were designed to develop a practical, quick and reliable screening system for basal stem rot (BSR) disease of oil palm using rubber wood inoculum block as an infection foci. Inoculation was performed under non-sterile conditions. 100% success on establishment of disease was achieved on seedlings and immature palms in a relatively short period. *G. boninense* was able to invade and colonise the oil palm roots without mechanical wounding.

Close contact between *G. boninense* and oil palm seedlings was established in a specially designed plastic container. Disease associated-symptoms were observed within two to three months after inoculation. The typical symptoms observed were the progressive yellowing and desiccation of leaves from the oldest to the youngest and the presence of basidioma. The extend of foliar symptoms reflected the disease severity of the roots and bole. The foliar symptoms were more severe when basidioma were detected. Mean severity of foliar symptoms was 42%, 33% and 16% on seedlings with fully developed basidioma, basidioma primodia and

without basidioma, respectively. Arrested growth and development were also observed on the seedlings. This was later followed by the death of the seedlings.

By placing the inoculum block in close contact with three roots and the bole of the one year old oil palm in the field, 15-25% infection of the bole tissues was achieved nine months after inoculation. The BSR associated symptoms observed were desiccation of the fronds and retardation of growth. BSR development tends to concentrate at the centre of the bole. Infected palms in the field were usually free from visible foliar symptoms until considerable damage had occurred in the bole region. Although all inoculated palms were infected, only 25% of the infected palms exhibited foliar symptoms 22 months after inoculation.

In vitro study showed that triazoles were significantly more effective than morpholines, with an EC_{50} values of 0.008-0.121 ppm a.i. and 1.939-2.944 ppm a.i., respectively. Total inhibition of mycelial growth by triazoles occurred at 1-5 ppm a.i. as compared to 250 ppm a.i. with morpholines. Difenconazole appeared to be the most active fungicide to the mycelial growth of *G. boninense*. However, the morpholines generally exhibited better vapour phase activity than the triazoles against mycelial growth of *Ganoderma*. Penconazole and tridemorph showed good vapour phase activity and uniform colony inhibition.

Uptake and efficacy of fungicides were detected by the bioassay on oil palm seedlings. Apparently penconazole had greater systemic activity than propiconazole and difenconazole. Fungicides also have phytotoxicity in the order penconazole > propiconazole > difenconazole. The greenhouse fungicide screening study on the bioefficacy indicated the potential curative activity of penconazole (0.5 g a.i., preliminary experiment) and both difenconazole and propiconazole (1.5 g a.i., repeat experiment) against BSR during early infection stage. However, activity decreased with increase in degree of colonisation, especially on seedlings which have already produced basidioma.

Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi syarat untuk Ijazah Master Sains Pertanian.

**PENYAKIT REPUT PANGKAL BATANG PADA KELAPA SAWIT:
SISTEM PENYARINGAN DAN KEBERKESANAN PERENCAT
STEROL BIOSINTESIS TERHADAP *Ganoderma boninense***

Oeh

TEH KIM SING

Mei 1996

Pengerusi : Profesor Madya Dr. Sariah Meon

Faculty : Pertanian

Kajian dijalankan untuk mendapatkan satu sistem penyaringan bagi penyakit reput pangkal batang (BSR) kelapa sawit yang praktikal, cepat dan berkesan dengan penggunaan inokulum blok kayu getah sebagai sumber jangkitan. Penginokulan dilakukan dalam keadaan yang tidak steril. Dalam tempoh yang singkat, 100% kejayaan jangkitan telah tercatat pada anak benih kelapa sawit dan kelapa sawit yang belum matang. Cederaan tidak perlu untuk memasukkan dan pengkolonian *G. boninense* pada akar kelapa sawit.

Sentuhan rapat antara *G. boninense* and anak benih kelapa sawit telah diwujudkan di dalam sebuah kotak plastik yang direka khas. Simptom yang berkaitan dengan penyakit dikesan dalam jangka masa dua hingga tiga bulan selepas inokulasi. Simptom yang biasa kelihatan adalah kekuningan dan pengeringan daun secara progresif bermula daripada daun peringkat tua ke muda dan kehadiran basidioma. Lanjutan simptom foliar memberi gambaran keterukan penyakit pada akar dan

pangkal batang pokok. Simptom tersebut bertambah teruk apabila terdapat basidioma. Anak benih yang memiliki basidioma, primodia basidioma dan tanpa basidioma masing-masing menunjukkan purata keterukan simptom foliar 42%, 33% dan 16%. Sekatan pertumbuhan dan perkembangan juga berlaku diikuti dengan kematian anak benih.

Dengan penempatan blok inokulum bersentuh rapat dengan tiga batang akar dan pangkal batang kelapa sawit yang berusia satu tahun, telah menghasilkan 15-25% jangkitan pada tisu-tisu pangkal batang sembilan bulan selepas penginokulan. Simptom berkaitan dengan penyakit BSR yang dikesan adalah pengeringan pelepah daun dan pertumbuhan yang bantut. Perkembangan BSR lebih tertumpu kepada bahagian tengah pangkal batang pokok. Di ladang, pokok yang terjangkit selalunya bebas daripada simptom foliar yang nyata sehingga kerosakkan menular ke dalam pangkal batang. Walaupun keseluruhan pohon yang diinokulat terjangkit, hanya 25% daripadanya menunjukkan simptom foliar 22 bulan selepas penginokulan.

Data *In vitro* menunjukkan bahawa triazole lebih berkesan secara bermakna daripada morpholine, dengan nilai EC_{50} 0.008-0.121 bsj b.a. dan 1.939-2.944 bsj b.a., masing-masing. Untuk triazole, perencatan pertumbuhan miselium sepenuhnya berlaku pada 1-5 bsj b.a. berbanding dengan morpholine pada 250 bsj b.a. Difenconazole merupakan racun kulat yang paling aktif terhadap pertumbuhan miselium *G. boninense*. Akan tetapi, morpholine pada umumnya menunjukkan aktiviti fasa wap yang lebih aktif daripada triazole merencat pertumbuhan miselium *Ganoderma*. Penconazole dan tridemorph menunjukkan aktiviti fasa wap yang baik dan perencatan koloni yang seragam.

Penyerapan dan keberkesanan racun kulat telah dikesan melalui bioasai anak benih kelapa sawit. Penconazole mempunyai aktiviti sistemik yang lebih tinggi daripada propiconazole dan difenconazole. Racun kulat juga menghasilkan

ketoksikan ke atas pokok dalam susunan penconazole > propiconazole > difenoconazole. Kajian keberkesanan racun kulat di rumah kaca menunjukkan pada peringkat awal jangkitan, penconazole (0.5 g b.a., eksperimen awalan) dan kedua-dua difenoconazole dan propiconazole (1.5 g b.a., eksperimen ulangan) berpotensi menghasilkan aktiviti pembaik pulih daripada penyakit BSR, tetapi aktiviti ini berkurangan dengan pertambahan tahap pengkolonian, terutama ke atas anak benih kelapa sawit yang telah menghasilkan basidioma.

CHAPTER I

INTRODUCTION

Oil palm (*Elaeis guineensis* Jacq.) was introduced to Malaysia in 1930. However, interest to cultivate and expand this world's most productive oil crop only picked up in the early 1960's. This change of interest was the result of the agricultural diversification programme implemented by the Malaysian Government. In a period of thirty years, the area under oil palm has increased twenty-fold to almost two million hectares. Palm oil is now Malaysia's major economic contributor to the agricultural sector. The mid to long-term price forecastings are stimulating the plantation groups to expand their oil palm area at the expense of rubber and cocoa. It is anticipated that the area under oil palm will further advance to 2.6 million hectares by the year 2000.

Recently the plantation sectors are facing numerous critical challenges. Areas of concern include shortage of human resources, cost of production, yield potential, rival oils and/or fats, competition from neighbouring countries and pest problems. Realising the importance of these potential predicaments, the Government and the plantation institutions are actively investigating feasible solutions. For instance, the Palm Oil Research Institute Malaysia (PORIM), Universiti Pertanian Malaysia (UPM) and Oil Palm Growers' Association have formed an Action Committee to address the problem of basal stem rot (BSR) of oil palm, a serious fungal malady in this region.

BSR of oil palm, caused by *Ganoderma* spp., is recognised to be one of the most important diseases of oil palm for sustainable economic production. This BSR

disease was initially thought to be unimportant, and frequently associated with old ageing palms in the coastal areas. Presently, this disease is frequently found in 10-15 year-old fields, especially those located around the coastal region of lower Perak and at sites previously planted with coconut (Singh, 1991; Khairudin, 1990; Turner, 1981). A further shift in disease distribution was recorded recently. Occurrence of BSR was found on young palms of the second or third generation and in peat and inland areas (Benjamin, 1993; Singh, 1991; Ariffin *et al.*, 1989a). Widya *et al.* (1991) reported that the incidence of BSR in Indonesia is more prominent on palms of the second and third generation.

The planting techniques and/or previous crop appear to have close relationship on the speed of BSR appearance and disease incidence (Chung *et al.*, 1994; Khairudin, 1994; Singh, 1991). Higher incidence of BSR on young plantings before reaching their most productive period is anticipated. This is alarming as between 1995-2000 about 0.4 million hectares of oil palm will be replanted and cultivated under large monoculture systems. Most of these plantings will be the second and/or third generation cultivation. Besides, the adoption of the cost-effective under-planting technique of oil palm, will further contribute to the spread of BSR disease in immature fields.

The potential occurrence of disease in immature fields certainly is a threat to the industry. Until now, early diagnosis of the disease is impossible. Confirmation of BSR is based on the presence of basidiomata. This is inadequate as usually immature palms seldom produce basidiomata, even at the advanced stage of disease development (Benjamin, 1993). However, when above ground symptoms are noticeable, extensive damage has already taken place at the palm base. This scenario can lead to large vacant fields as currently experienced by some of the coastal and/or riverine estates.

Currently, there is no effective control of BSR palm in the field. The existing practice of the industry emphasising on preventive measure through clean clearing.



Complete removal of the infected tissues and root masses is impossible in the field, thus, the immature palms remain at risk to BSR. Besides, *Ganoderma* appears to produce resting structures in infected woody remains (Ariffin *et al.*, 1989b).

There is an urgent need to control BSR. Nevertheless, to conduct field trials on this disease is not an easy task. Critical constraints include the selection of uniform trial plots, slow progress of disease and inconsistency of trial methodology are still existing. It is extremely important to have a system where a preliminary test can be concluded in a short period under semi-control condition. Hence, the present studies were conducted with the following objectives:

- To develop a fast and reliable screening system of BSR on seedlings and immature oil palms using a rubber wood inoculum block
- To evaluate *in vitro* bioefficacy of sterol-biosynthesis inhibitors on *G. boninense* with direct contact and their vapour phase activity
- To investigate the *in vivo* behaviour and curative activity of triazoles against BSR on inoculated oil palm seedlings

CHAPTER II

LITERATURE REVIEW

Status of *Ganoderma* spp. as a Pathogen of Basal Stem Rot (BSR) of Oil Palm

Basal stem rot of oil palm was first reported in Malaya towards the end of 1927 and was initially associated with termites attack. Later, bracket-like fructifications emerged and were identified as *G. lucidum* (Leys) Karst (Thompson, 1931). In 1935, Thompson considered *G. lucidum* as pathogen of oil palms causing stem rot mainly of leaning palms (Thompson, 1935). It was described as basal stem rot by Voelcker (1950-51). BSR of oil palm was regarded a pathogen of some important when a patch of palms over 20 year-old was found infected (Heath, 1956). Initially, this malady did not arouse immediate attention since it had been viewed as an old age problem. Wijbrans (1955) considered that *G. lucidum* was a facultative parasite which would cause damage only on palms weakened by injuries and unfavourable environmental conditions. Turner believed that the old palms were susceptible because of the break-down in immunity barriers (Turner, 1981). Incidence of BSR on eight-year-old palms and subsequently on younger palms were observed for the first time in 1957 (Heath, 1957).

Several reports on BSR distribution in the country indicated close association of the disease incidence with areas previously planted with coconut (Turner, 1965a and 1965b; Navaratnam, 1964). There were signs that stumps left in the ground behaved as infection foci. In the 1980's, large hectareage of oil palm infected with BSR, particularly in the coastal areas and at sites formerly planted with coconut were reported. The disease reached endemic levels on 10 to 15-year-old plantings (Singh, 1991; Turner, 1981). Of late, the incidence of BSR has been

recorded on peat areas, inland areas and on young palms of the second and/or third generation (Benjamin, 1993; Singh, 1991; Ariffin *et al.*, 1989a). The fungus colonised young palms in a short period of time causing toppling over and/or die-standing of palms, even before reaching the optimum production stage. This leads to substantial economic loss which are due to low stand per unit cultivated area and reduced yield on diseased palms (Khairudin, 1994 and Singh, 1991).

Although much is known about the incidence of BSR, the confusion with regards to the identity of the pathogen has yet to be solved. In Malaysia, it was first identified as *G. lucidum* Karst by Thompson (1931). Varghese (1965) and Turner and Bull (1967) suggested that a complex of species might be involved. Identification made by Steyaert (1967) on morphological characteristics of sporophores indicated that more than one species of *Ganoderma* were associated with BSR in Malaysia. Based on the association with BSR from various parts of the world, Turner (1981) recorded at least 15 species of *Ganoderma* as likely pathogens of oil palm. However, based on associations and morphological studies of basidioma collected from various location in Peninsular Malaysia, Ho and Nawawi (1985) concluded that *G. boninense* was the sole pathogen, however, possible differential pathogenicity of the fungus was not confirmed. Thus, numerous problems on species or strains are yet to be resolved.

In the nineties, a significant progress to confirm the pathogenicity of *Ganoderma* was attained. With the development of a reliable inoculation technique (Ariffin *et al.*, 1995; Sariah *et al.*, 1994 and Khairudin *et al.*, 1991) consistently high rates of BSR establishment on oil palm seedlings were achieved. Khairudin (1991) also showed that *G. boninense* was pathogenic to oil palm, whilst *G. philippi* and *G. lucidum* were not. Ariffin *et al.* (1991a) using selective media consistently isolated *G. boninense* from diseased palms, with and without visual symptoms. *G. boninense* is now established as the causal agent of BSR.

Infection of *G. boninense* takes place via root contact between healthy palms and the inoculum sources in the field, particularly infected dead stump tissues. The pathogen then colonises the roots and spread to the other tissues. Hitherto, the role of basidiospores of *G. boninense* in disease development is not clear. Inoculation of spores onto the cut frond bases (Turner, 1965c) and oil palm seedlings failed to yield infection (Ramasamy, 1972; Yeong, 1972). Turner (1991) believed that spores are responsible for spread into new planting areas. The possible role of insects in transmitting spores to healthy palms could not be ruled out.

Symptoms of Basal Stem Rot on Oil Palm

Foliar Symptoms

Inoculated Seedlings

Typical foliar symptoms of infection observed are progressive yellowing and desiccation of leaves from the oldest to the youngest. Desiccation of a leaf usually starts at the tip followed by rapid yellowing and drying of the entire lamina (Khairudin, 1990). Symptoms of yellowing and desiccation of leaves only appear in cases where infection has reached the bole, and generally appear on the same side of the inoculated root (Ariffin *et al.*, 1995). The extent of foliar symptoms is not indicative of the damage to the roots and bole region (Sariah *et al.*, 1994). Contrary to this, Khairudin (1990) reported that the severity of rot in the roots and/or bole was reflected by the foliar symptoms. Seedlings showed arrested growth and development, and later foliage dried-up and death of seedlings occurred (Khairudin, 1990).

Immature Palms

A typical early symptom is the occurrence of frond marginal necrosis starting from the lower whorl and later extending upwards. This is followed by chlorosis or mottling and desiccation. One-sided foliar symptoms are most common (Benjamin, 1993; Singh, 1991). This may serve as an indication of the site of infection. The

newly unfolded leaves are shorter and chlorotic, and the tips may be necrotic. As the disease progresses, the palms take on a pale appearance, produce a number of unopened spears and growth becomes retarded (Singh, 1991).

Mature Palms

The symptoms resemble palms suffering from water deficiency. Excessive production of unopened spears are common. Subsequently, desiccation and wilting of fronds occurs. It usually begins with the oldest fronds and extends progressively to the younger ones and later to the crown. General nutrient deficiency symptoms may also appear. Desiccated fronds droop or break at the rachis and encircle the trunk. Often, when foliar symptoms are observed, more than 50% of the cross-sectional area of the palm base is necrotic. Eventually, palms die standing, and may remain erect or topple-over (Khairudin, 1994; Turner, 1981).

Basidioma

Inoculated Seedlings

Around the bole region, profuse vegetative growth of white masses rhizomorph can occur three to five weeks after inoculation, and later develop into a cushion-like stroma. This mycelia cushion rapidly becomes button-like (basidioma primodia) and is later followed by the apparent development of stipe. Primodia of basidioma develop into bracket shape basidioma within two weeks (Khairudin, 1990). Sariah *et al.* (1994) reported that typical white margin, bracket shape basidioma are formed within 12 to 16 weeks after inoculation. The position of the basidioma indicates the rough position of diseased area in the bole.

Immature and Mature Palms

On mature palms, basidioma are produced as the decay advances and usually develop at the base of the stem or on diseased primary roots. They may form before and/or after foliar symptoms. They begin as white basidioma primodia, but soon expand and become bracket-like. The mature basidioma vary in shape, size and colour. The shiny lacquered finish upper surface can be light to dark red brown, with

a swollen white margin. The lower surface is greyish white in colour and has numerous minute pores. Basidioma are seldom seen on infected young palms. As the bole tissue of young palm is soft, it decays rapidly and palms usually die before the fruiting-bodies are produced (Turner, 1966; Singh, 1991).

Symptoms in the Roots and Stem Region (Bole)

Inoculated Seedlings

Severe root rotting has been reported in seedlings which had displayed progressive desiccation of the foliage. The cortex and epidermis of infected roots were readily separated from the stele (Khairudin, 1990). The fungus was detected on all root tissues, but was more pronounced in the vascular system (Sariah *et al.*, 1994). Ariffin *et al.* (1995) when screening for resistance revealed that the average movement of *G. boninense* within infected roots for all progenies tested was 10.22 mm/month.

On diseased tissues of the bole, progressive lesions were noted which later extended into the stem region as the fungus colonised further (Sariah *et al.*, 1994).

Immature and Mature Palms

In infected palms, the diseased roots are friable and brittle. The cortex becomes brown and the stele black. In the older roots, the fungus may be present as a whitish skin-like layer on the inner surface of the exodermis. The dry rotten tissues of the bole and stem tissues appear light brown. Interspersed within the decaying tissues are narrow dark lines or bands, which contain chlamyospore-like structures (Ariffin *et al.*, 1989b). Immediately above the advancing margin of the lesion, a lemon-yellow zone, presumably the result of host-parasite reaction is present (Singh, 1991; Turner, 1981). White mycelial of the fungus can be seen inside numerous small cavities of the light brown diseased tissues (Turner, 1981).

Establishment of *Ganoderma* Infection by Inoculation

Several techniques were used to establish the pathogenicity of *Ganoderma* on oil palm. Navaratnam (1961) carried out inoculation with a pure culture of *Ganoderma lucidum* (grew on sterilised sand and maize-meal medium) on the roots and stem tissues of old palms. The rate of success infection was 5% and 15% for root and stem inoculation, respectively. He ascribed the limited success rate to faulty technique and poor establishment of inoculum. Nevertheless, this observation has revealed that both oil palm root and stem were susceptible to *G. lucidum* infection.

Subsequently, successful inoculations of palm seedlings using large infected tissues were also reported (Navaratnam and Chee, 1965; Peries, 1974). However, Ramasamy (1972) and Ariffin *et al.* (1993) failed to obtain infection with this method. Inoculation of spores to different parts of oil palm also failed to yield infection (Turner, 1965c; Ramasamy, 1972; Yeong, 1972). Turner (1981) suggested that spores have inadequate inoculum potential for direct infection of a living palm.

Success of inoculation of *Ganoderma* was also reported on other crops. Adaskaveg and Gilbertson (1987) obtained a high success rate of infection by placing *G. lucidum* infected wooden blocks adjacent to wounded or unwounded roots of grapevines. Further inoculation on trunk tissues of field grapevine with infected grape wood chips was also successful. Sampath Kumar and Nambiar (1990) obtained positive pathogenicity of *Ganoderma spp.* on arecanut using stem insertion methods and planting of disease stump around healthy stem of arecanut.

Recently, a significant progress on disease establishment with pure culture of *G. boninense* on oil palm seedlings was demonstrated. Apparently, there are some similarity among the approaches. Firstly, pure cultures of *G. boninense* produced in the laboratory were used as inoculum. Secondly, inocula used were in a vigorous state and thirdly, close contact between the pathogen and host was made.

Okky *et al.* (1990) inoculated oil palm plantlets and calli *in vitro* with *G. boninense*. This technique had provided a contamination free environment to initiate the host-pathogen relationship. The mortality rate of the plantlets from those with wounded basal stems, wounded roots and unwounded plantlets were 90%, 35% and 45% respectively. They concluded that inoculation on wounded basal stem of plantlet was the best method.

Khairudin (1990) used mesocarp fibre of oil palm and rubber wood blocks as a substrate to culture *G. boninense* for the pathogenicity test on oil palm seedlings. Three to four leaf stage seedlings were placed on five-week-old inoculum in polypropylene bags and covered with sterilised soil. At nine months after inoculation, only two out of the 18 inoculated seedlings of the mesocarp fibre inoculum were infected. In the case where *G. boninense* was cultured on rubber wood blocks and placed in close contact with the roots of oil palm seedlings, excellent success infection rate was reported. Foliar symptoms were noticeable at seven months after inoculation, and by 11 months after inoculation all inoculated seedlings were infected with two dead. He concluded that the rubber wood block is an excellent substrate for the establishment of *G. boninense* inoculum. He further described that rubber wood, being solid, would enhanced the survival of the pathogen. Subsequent studies of Khairudin (1994) and Sariah *et al.* (1994) with rubber wood inoculum block had repeatedly yielded consistent and high success rates on oil palm seedlings. There appear to be a positive correlation existed between the size of inoculum and the degree of disease incidence (Khairudin, 1994). At 11 months, the incidence of disease for the 54, 108, 216 and 432 cm³ rubber wood block inoculum were 5.9%, 5.9%, 17.6% and 82.4%, respectively.

Ariffin and Idris (1993) reported that their earlier attempts to artificially inoculate oil palm seedlings and field palms with pure cultures of *G. boninense* were unsuccessful due to contamination. Therefore, they later developed a single root inoculation technique with pure culture of *G. boninense*. During inoculation, a primary root was selected, surface sterilised, and then the distal end of the root was